

Variational Methods & Optimal Control

lecture 02

Matthew Roughan

<matthew.roughan@adelaide.edu.au>

Discipline of Applied Mathematics
School of Mathematical Sciences
University of Adelaide

April 14, 2016

Variational Methods & Optimal Control: lecture 02 – p.1/29

Revision, part ii


Extrema of functions of multiple variables. Taylor's theorem and the chain rule in N-D. Hessians and classification of extrema.

Variational Methods & Optimal Control: lecture 02 – p.2/29

Extrema of functions of one variable

Local extrema have $f'(x) = 0$

includes maxima, minima, and stationary points of inflection




Variational Methods & Optimal Control: lecture 02 – p.3/29

Classification of extrema

Local extrema have $f'(x) = 0$

- ▶ $f''(x) > 0$ local minima
- ▶ $f''(x) < 0$ local maxima
- ▶ $f''(x) = 0$ it might be a stationary point of inflection, depending on higher order derivatives, e.g. x^4 .



Variational Methods & Optimal Control: lecture 02 – p.4/29

Functions of n variables

- ▶ Let Ω be a closed region of \mathbb{R}^n , i.e. $\Omega \subset \mathbb{R}^n$
- ▶ Let $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \Omega$
- ▶ Let $f : \Omega \rightarrow \mathbb{R}$
- ▶ A local minima if $f(\cdot)$ is point \mathbf{x} such that there exists $\delta > 0$ where

$$f(\hat{\mathbf{x}}) \geq f(\mathbf{x})$$

for any $\hat{\mathbf{x}} \in B(\mathbf{x}; \delta)$.


- ▶ A global minima of $f(\cdot)$ on Ω is point \mathbf{x} such that

$$f(\hat{\mathbf{x}}) \geq f(\mathbf{x})$$

for any $\hat{\mathbf{x}} \in \Omega$.


2D example 2

- ▶ $f(x_1, x_2) = r - 1/2r^2$, where
 $r = \sqrt{x_1^2 + x_2^2}$
- ▶ global maxima on curve $r = 1$
- ▶ local minima at $r = 0$




2D example 1

- ▶ $f(x_1, x_2) = x_1^2 - x_2^2 + x_1^3$
- ▶ local maximum at $(-2/3, 0)$
- ▶ saddle point at $(0, 0)$



2D example 3

- ▶ $f(x_1, x_2) = x_2^3 - 3x_1^2x_2$
- ▶ Monkey saddle at $(0, 0)$



The Chain rule

The derivative of a function $f(x_1, x_2)$ along a line described parametrically by $(x_1(t), x_2(t))$

$$\frac{df}{dt} = \frac{\partial f}{\partial x_1} \frac{dx_1}{dt} + \frac{\partial f}{\partial x_2} \frac{dx_2}{dt}$$

Another way to think of this is as the directional derivative formed from the dot product of grad and the direction of the line, e.g.,


$$\begin{aligned} \frac{df}{dt} &= \nabla f \cdot \frac{d\mathbf{x}}{dt} \\ &= \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2} \right) \cdot \left(\frac{dx_1}{dt}, \frac{dx_2}{dt} \right) \end{aligned}$$

A graphical example

For a function of two variables $f(x, y)$ we get

$$\frac{df}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}$$

$$\begin{aligned} f(x, y) &= x + y \\ x &= \cos t \\ y &= \sin t \end{aligned}$$



Chain Rule for more variables

The chain rule (for a function of more than one variable)

$f(\mathbf{x}) = f(x_1, x_2, \dots, x_n)$, where we want to find the derivative of a function $f(\mathbf{x})$ along a line described parametrically by $(x_1(t), x_2(t), \dots, x_n(t))$ then we take

$$\frac{df}{dt} = \frac{\partial f}{\partial x_1} \frac{dx_1}{dt} + \frac{\partial f}{\partial x_2} \frac{dx_2}{dt} + \dots + \frac{\partial f}{\partial x_n} \frac{dx_n}{dt}$$

or alternatively

$$\begin{aligned} \frac{df}{dt} &= \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right) \cdot \left(\frac{dx_1}{dt}, \frac{dx_2}{dt}, \dots, \frac{dx_n}{dt} \right) \\ &= \nabla f \cdot \frac{d\mathbf{x}}{dt} \end{aligned}$$

Chain Rule Derivation

By the definition

$$\frac{df}{dt} = \lim_{\epsilon \rightarrow 0} \frac{f(x(t+\epsilon), y(t+\epsilon)) - f(x(t), y(t))}{\epsilon}$$

But note that from Taylor's theorem

$$x(t+\epsilon) = x(t) + \epsilon x'(t) + O(\epsilon^2).$$

As we consider the limit as $\epsilon \rightarrow 0$ we may ignore the $O(\epsilon^2)$ term, to get

$$\frac{df}{dt} = \lim_{\epsilon \rightarrow 0} \frac{f(x(t) + \epsilon x'(t), y(t) + \epsilon y'(t)) - f(x(t), y(t))}{\epsilon}$$

Chain Rule Derivation

$$\begin{aligned}
 \frac{df}{dt} &= \lim_{\varepsilon \rightarrow 0} \frac{f(x(t) + \varepsilon x'(t), y(t) + \varepsilon y'(t)) - f(x(t), y(t))}{\varepsilon} \\
 &= \lim_{\varepsilon \rightarrow 0} \frac{f(x(t) + \varepsilon x'(t), y(t) + \varepsilon y'(t)) - f(x(t), y(t) + \varepsilon y'(t))}{\varepsilon} \\
 &\quad + \lim_{\varepsilon \rightarrow 0} \frac{f(x(t), y(t) + \varepsilon y'(t)) - f(x(t), y(t))}{\varepsilon} \\
 &= x'(t) \lim_{\varepsilon \rightarrow 0} \frac{f(x(t) + \varepsilon x'(t), y(t) + \varepsilon y'(t)) - f(x(t), y(t) + \varepsilon y'(t))}{\varepsilon x'(t)} \\
 &\quad + y'(t) \lim_{\varepsilon \rightarrow 0} \frac{f(x(t), y(t) + \varepsilon y'(t)) - f(x(t), y(t))}{\varepsilon y'(t)}
 \end{aligned}$$

Variational Methods & Optimal Control: lecture 02 – p.13/29

Chain Rule Derivation

$$\begin{aligned}
 \frac{df}{dt} &= x'(t) \lim_{\varepsilon \rightarrow 0} \frac{f(x(t) + \varepsilon x'(t), y(t) + \varepsilon y'(t)) - f(x(t), y(t) + \varepsilon y'(t))}{\varepsilon x'(t)} \\
 &\quad + y'(t) \lim_{\varepsilon \rightarrow 0} \frac{f(x(t), y(t) + \varepsilon y'(t)) - f(x(t), y(t))}{\varepsilon y'(t)} \\
 &= x'(t) \lim_{\varepsilon_x \rightarrow 0} \frac{f(x(t) + \varepsilon_x, y(t) + \varepsilon y'(t)) - f(x(t), y(t) + \varepsilon y'(t))}{\varepsilon_x} \\
 &\quad + y'(t) \lim_{\varepsilon_y \rightarrow 0} \frac{f(x(t), y(t) + \varepsilon_y) - f(x(t), y(t))}{\varepsilon_y} \\
 &= x'(t) \frac{\partial f}{\partial x} + y'(t) \frac{\partial f}{\partial y}
 \end{aligned}$$

which is the chain rule!

Variational Methods & Optimal Control: lecture 02 – p.14/29

Chain Rule special case

When we only have one variable, we simply want to calculate the derivative of a function f of another function x , e.g.

$$\frac{d}{dt} f(x(t)) = \frac{df}{dx} \frac{dx}{dt}$$

Another way of writing this is

$$\frac{d}{dt} f(x(t)) = f'[x(t)] x'(t),$$

which is the form you learnt in 1st year.

Variational Methods & Optimal Control: lecture 02 – p.15/29

Taylor's theorem in 2D

$$\begin{aligned}
 f(x_1 + \delta x_1, x_2 + \delta x_2) &= f(x_1, x_2) + \delta x_1 \frac{\partial f}{\partial x_1} + \delta x_2 \frac{\partial f}{\partial x_2} \\
 &\quad + \frac{1}{2} \left[\delta x_1^2 \frac{\partial^2 f}{\partial x_1^2} + 2\delta x_1 \delta x_2 \frac{\partial^2 f}{\partial x_1 \partial x_2} + \delta x_2^2 \frac{\partial^2 f}{\partial x_2^2} \right] + \dots
 \end{aligned}$$

Write $(\delta x_1, \delta x_2) = \varepsilon \times (\eta_1, \eta_2)$

$$\begin{aligned}
 f(\mathbf{x} + \varepsilon \boldsymbol{\eta}) &= f(\mathbf{x}) + \varepsilon \left(\eta_1 \frac{\partial f}{\partial x_1} + \eta_2 \frac{\partial f}{\partial x_2} \right) \\
 &\quad + \frac{\varepsilon^2}{2} \left[\eta_1^2 \frac{\partial^2 f}{\partial x_1^2} + 2\eta_1 \eta_2 \frac{\partial^2 f}{\partial x_1 \partial x_2} + \eta_2^2 \frac{\partial^2 f}{\partial x_2^2} \right] + O(\varepsilon^3)
 \end{aligned}$$

Variational Methods & Optimal Control: lecture 02 – p.16/29

Taylor's theorem in N-D

$$f(\mathbf{x} + \delta\mathbf{x}) = f(\mathbf{x}) + \sum_{i=1}^n \delta x_i \frac{\partial f}{\partial x_i} + \frac{1}{2} \sum_{i,j=1}^n \frac{\partial^2 f}{\partial x_i \partial x_j} \delta x_i \delta x_j + O(\delta\mathbf{x}^3)$$

$$f(\mathbf{x} + \delta\mathbf{x}) = f(\mathbf{x}) + \delta\mathbf{x}^T \nabla f(\mathbf{x}) + \frac{1}{2} \delta\mathbf{x}^T H(\mathbf{x}) \delta\mathbf{x} + O(\delta\mathbf{x}^3)$$

Where $H(\mathbf{x})$ is the Hessian matrix

$$H(\mathbf{x}) = \begin{pmatrix} \frac{\partial^2 f}{\partial x_1^2} & \frac{\partial^2 f}{\partial x_1 \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n} \\ \frac{\partial^2 f}{\partial x_2 \partial x_1} & \frac{\partial^2 f}{\partial x_2^2} & \cdots & \frac{\partial^2 f}{\partial x_2 \partial x_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1} & \frac{\partial^2 f}{\partial x_n \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_n^2} \end{pmatrix}$$

Maxima of N variables

If a smooth function $f(\mathbf{x})$ has a local extrema at \mathbf{x} then

$$\nabla f(\mathbf{x}) = \left(\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_n} \right)^T = \mathbf{0}$$

A sufficient condition for the extrema \mathbf{x} to be a local minimum is for the quadratic form

$$Q(\delta x_1, \dots, \delta x_n) = \delta\mathbf{x}^T H(\mathbf{x}) \delta\mathbf{x} = \sum_{i=1}^n \sum_{j=1}^n \frac{\partial^2 f}{\partial x_i \partial x_j} \delta x_i \delta x_j$$

to be strictly positive definite.

Quadratic forms

A quadratic form

$$Q(\mathbf{x}) = \sum_{i,j} a_{ij} x_i x_j = \mathbf{x}^T A \mathbf{x}$$

is said to be positive definite if $Q(\mathbf{x}) > 0$ for all $\mathbf{x} \neq \mathbf{0}$.

A quadratic form is positive definite iff every eigenvalue of A is greater than zero.

A quadratic form is positive definite if all the principal minors in the top-left corner of A are positive, in other words

$$a_{11} > 0, \quad \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} > 0, \quad \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} > 0, \dots$$

Notes on maxima and minima

- ▶ maxima of $f(x)$ are minima of $-f(x)$.
- ▶ haven't said anything about non-differentiable functions
- ▶ if continuous in the interval, must achieve maximum (minimum) in the interval

Calculus of variations

- ▶ We are not maximizing the value of a function.
- ▶ We are maximizing a **functional** — a function of a function
- ▶ Can think of it as an ∞ -dimensional max. problem.
 - ▷ can choose between different functions
 - ▷ function sits in ∞ -dimensional vector space
- ▶ This might take some effort.

Integral functionals

- ▶ Previous functionals not very interesting.
- ▶ Easy to find $y(x)$ which minimizes these.
- ▶ Integral functionals are more interesting.
- ▶ Example integral functionals

$$F\{y\} = \int_a^b y(x) dx$$

$$F\{y\} = \int_a^b f(x)y(x) dx$$

$$F\{y\} = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

Functionals

A **Functional** maps an element of a vector space (e.g. a space containing functions) to a real number, e.g. $F : S \rightarrow \mathbb{R}$.

Example Functionals

$$F\{y(x)\} = |y(0)|$$

$$F\{y(x)\} = \max_x \{y(x)\}$$

$$F\{y(x)\} = \left. \frac{dy}{dx} \right|_{x=1}$$

$$F\{y(x)\} = y(0) + y(1)$$

$$F\{y(x)\} = \sum_{n=0}^N a_n y(n)$$

Some simple cases

$$F\{y\} = \int_{a(\epsilon)}^{b(\epsilon)} y(x, \epsilon) dx$$

$$\frac{dF}{d\epsilon} = y(b, \epsilon) \frac{db}{d\epsilon} - y(a, \epsilon) \frac{da}{d\epsilon} + \int_{a(\epsilon)}^{b(\epsilon)} \frac{\partial y(x, \epsilon)}{\partial \epsilon} dx$$

If a and b are fixed then

$$\frac{da}{d\epsilon} = 0$$

$$\frac{db}{d\epsilon} = 0$$

and so the derivative of the integral becomes the integral of the derivative.

Crude Brachystochrone

Brachystochrone involves the functional

$$F\{y\} = \int_{x_0}^{x_1} \sqrt{\frac{1+y^2}{y}} dx$$


Let us guess that the brachystochrone takes the form

$$y(x, \epsilon) = (1-x)^\epsilon$$

We could calculate the derivative WRT ϵ as above and compute the stationary points by finding


$$\frac{dF}{d\epsilon} = 0$$

Crude Brachystochrone



► but what if the family of curves doesn't contain the maximum?

Crude Brachystochrone



Extra bits

Notation

- ▶ $f(\mathbf{x}) : S \rightarrow \mathbb{R}$ denotes a function that maps the set $S \subset \mathbb{R}^n$ to a real number.
- ▶ $\frac{\partial^n f}{\partial x_i^n}$ denotes the n th partial derivative of $f(\mathbf{x})$, with respect to x_i .
- ▶ the ε -neighborhood under the Euclidean norm is $B(\mathbf{x}; \varepsilon) = \{ \hat{\mathbf{x}} \in \mathbb{R}^n \mid \|\hat{\mathbf{x}} - \mathbf{x}\|_2 < \varepsilon \}$
- ▶ The Euclidean norm in \mathbb{R}^n is $\|\mathbf{x}\|_2 = \sqrt{\sum_{i=1}^n x_i^2}$
- ▶ $F\{y\}$ denotes a functional of the function $y(x)$.